Developmental Cell

Co-option of an Ancestral Hox-Regulated Network Underlies a Recently Evolved Morphological Novelty

Graphical Abstract



Highlights

- Newly evolved Drosophila adult genital structure allows analysis of network history
- The adult structure evolved by co-opting the network of a larval breathing structure
- Ten genes, including seven transcription factors, are shared between both networks
- Seven embryonic regulatory sequences are re-deployed during genital development

Authors

William J. Glassford, Winslow C. Johnson, Natalie R. Dall, ..., Werner Boll, Markus Noll, Mark Rebeiz

Correspondence rebeiz@pitt.edu

In Brief

Network reuse, or co-option, has been implicated in explaining the origins of many complex structures. Glassford et al. demonstrate that a newly evolved morphological structure in adult genitalia that is unique to the Drosophila melanogaster subgroup co-opted an embryonic transcriptional network that contributes to the development of a larval breathing structure.





Co-option of an Ancestral Hox-Regulated Network Underlies a Recently Evolved Morphological Novelty

William J. Glassford,¹ Winslow C. Johnson,¹ Natalie R. Dall,¹ Sarah Jacquelyn Smith,¹ Yang Liu,¹ Werner Boll,² Markus Noll,² and Mark Rebeiz^{1,*}

¹Department of Biological Sciences, University of Pittsburgh, 4249 Fifth Avenue, Pittsburgh, PA 15260, USA

²Institute of Molecular Life Sciences, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland

*Correspondence: rebeiz@pitt.edu

http://dx.doi.org/10.1016/j.devcel.2015.08.005

SUMMARY

The evolutionary origins of complex morphological structures such as the vertebrate eye or insect wing remain one of the greatest mysteries of biology. Recent comparative studies of gene expression imply that new structures are not built from scratch, but rather form by co-opting preexisting gene networks. A key prediction of this model is that upstream factors within the network will activate their preexisting targets (i.e., enhancers) to form novel anatomies. Here, we show how a recently derived morphological novelty present in the genitalia of D. melanogaster employs an ancestral Hox-regulated network deployed in the embryo to generate the larval posterior spiracle. We demonstrate how transcriptional enhancers and constituent transcription factor binding sites are used in both ancestral and novel contexts. These results illustrate network co-option at the level of individual connections between regulatory genes and highlight how morphological novelty may originate through the co-option of networks controlling seemingly unrelated structures.

INTRODUCTION

"...structural genes are building stones which can be used over again for achieving different styles of architecture...evolution is mostly the reutilization of essentially constituted genomes."

-Emile Zuckerkandl (Zuckerkandl, 1976)

Evolutionary biologists have long been intrigued by the origins of biological complexity. While the complexity of living systems can be considered at multiple levels of organization (e.g., the origins of DNA-based life [Crick, 1968; Orgel, 1968], organelles [Sagan, 1967], or multicellularity [Bonner, 1998]), the evolutionary origin of morphological complexity is a developmental problem (Muller and Wagner, 1991). Morphological structures are patterned and formed during the process of embryonic development, and each cell in the developing organism must derive unique physical properties from an identical DNA code. This apparent paradox is solved by differential gene activity, governed by vast gene regulatory networks (GRNs) (Davidson, 2001). Regulatory factors within GRNs bind transcriptional regulatory sequences such as enhancers to combinatorially determine the expression status of each gene of the network in morphological space and developmental time (Small et al., 1992). Hence, an understanding of the origins of morphological complexity necessitates investigations into how GRNs originate.

A growing body of evidence has implicated the re-use, or co-option, of existing networks in the evolution of novel morphological structures (Gao and Davidson, 2008; Keys et al., 1999; Kuraku et al., 2005; Moczek and Nagy, 2005). For example, expression of the appendage-patterning network within the developing beetle horn suggests that this novelty arose through the establishment of a new proximo-distal axis (Moczek and Nagy, 2005; Moczek and Rose, 2009; Moczek et al., 2006). Such findings evoke a scenario in which a cohort of downstream appendage enhancers was in turn activated in the new setting, generating a unique developmental output. However, instances of co-option have traditionally been supported by correlations in gene expression, relationships that may arise without the reuse of existing circuits (Abouheif, 1999). Currently, examples that illustrate this phenomenon at the level of enhancers and the constituent binding sites that were co-opted are lacking.

Here, we trace the evolutionary history of the posterior lobe, a recently evolved morphological structure present in the model organism *Drosophila* (*D.*) *melanogaster* at the level of its network, enhancers, and the transcription factor binding sites of which these are composed.

RESULTS

The Posterior Lobe Is a Morphological Novelty Unique to the *D. melanogaster* Subgroup

Male genitalia represent the most rapidly evolving morphological structures in the animal kingdom (Eberhard, 1985) and are often used to taxonomically distinguish insect species. The posterior lobe is a hook-shaped outgrowth unique to the external genitalia of *D. melanogaster* and its closest relatives in the *melanogaster* clade (Figure 1) (Jagadeeshan and Singh, 2006; Kopp and True, 2002). A cuticular projection similar to the posterior lobe is also present in the *yakuba* clade (Yassin and Orgogozo, 2013), suggesting a recent origin of this structure in the *melanogaster* subgroup (Figure S1). Among members of the *melanogaster* clade, the posterior lobe is highly divergent in shape





Figure 1. The Posterior Lobe Is a Morphological Novelty Unique to the *D. melanogaster* Clade

(A) Scanning electron micrograph of a *D. simulans* male with relevant structures labeled.

(B) Tree depicting the phylogenetic relationships of the species in this study, and brightfield images of their lateral plate cuticle morphologies. The posterior lobe is an outgrowth of the lateral plate unique to the *melanogaster* clade (arrows).

See also Figure S1.

and size and represents the only reliable character to distinguish species identity (Coyne, 1993). During mating, the posterior lobe is used by the male to grasp the female ovipositor (Jagadeeshan and Singh, 2006) and subsequently is inserted between cuticular plates at the posterior of the female abdomen during genital coupling (Robertson, 1988). Given the recent evolution of the posterior lobe and its presence in *D. melanogaster*, a highly tractable model organism for studying the structure and evolution of GRNs, we sought to elucidate its evolutionary origins.

An Ancestral Enhancer of *Pox neuro* Was Co-opted into the Posterior Lobe Network

To trace the evolutionary history of the posterior lobe, we first examined Pox neuro (Poxn), a gene that is critical to its development. Poxn encodes a paired-domain transcription factor required for proper posterior lobe formation (Boll and Noll, 2002). In a comprehensive survey of the regulatory region of Poxn, a segment spanning the second exon and intron (Figure 2A) was found to be required for posterior lobe development (Boll and Noll, 2002). To examine the role of this enhancer in genital development and identify how this role evolved, we cloned this segment of the D. melanogaster Poxn gene into a GFP reporter construct (Figure 2A). Transgenic animals bearing the genital enhancer of Poxn drive expression both before and during posterior lobe development. At 32 hr after puparium formation (hAPF), a time that precedes the formation of the posterior lobe (see Figures S2A–S2L for a time course of genital development in lobed and non-lobed species), we observed broad GFP expression in a zone that straddles the presumptive clasper and lateral plate (Figure 2D). As the posterior lobe emerges from the lateral plate and assumes its adult morphology, the reporter expresses high levels of GFP in the developing lobe (Figure 2E, arrow). This portion of the *Poxn* regulatory region accurately recapitulates the endogenous expression of *Poxn* mRNA and protein in the *D. melanogaster* lateral plate (Figures 2B and 2C; Figures S2M and S2N).

The high level of reporter and *Poxn* mRNA in the developing posterior lobe strongly suggests that *Poxn* plays a direct role during posterior lobe development. To examine how this role evolved, we first analyzed its expression in species that lack this structure. At 32 hAPF, the early pattern of *Poxn* expression in the non-lobed species *D. ananassae* greatly resembles that of *D. melanogaster* prior to posterior lobe formation (Figure 2F). However, *Poxn* expression quickly subsides in the *D. ananassae* lateral plate once it has separated from the clasper (Figure 2G). Similar results were obtained for two additional non-lobed species, *D. biarmipes* and *D. pseudoobscura* (Figures S2M and S2N), suggesting that late, high levels of *Poxn* expression are uniquely associated with the development of this novelty.

Differences in Poxn expression between lobed and non-lobed species may be due to changes in the posterior lobe enhancer region (i.e., in cis) or could be caused by changes in trans that altered upstream regulators in the genitalia (Wittkopp, 2005). To distinguish between these possibilities and ascertain whether the posterior lobe enhancer of Poxn recently derived its function, we examined the activity of this enhancer from species that lack this structure. Sequences orthologous to the D. melanogaster posterior lobe enhancer region were cloned from several non-lobed species and tested for the ability to drive GFP reporter expression in the D. melanogaster posterior lobe. The posterior lobe enhancer regions of D. ananassae, D. yakuba, and D. pseudoobscura Poxn all drove GFP expression that closely matched the pattern and timing of the D. melanogaster reporter construct (Figures 2H and 2I; Figures S3G'-S3I'). The ability of the posterior lobe enhancer region to produce strong expression in the developing posterior lobe despite the lack of this structure in these species strongly indicated that it predated the evolution of this novelty.

As our findings implied the absence of functionally significant changes in the *Poxn* enhancer during the evolution of the posterior lobe, we next tested whether a non-lobed species' enhancer could rescue the posterior lobe of a *D. melanogaster Poxn* mutant. The *D. melanogaster* posterior lobe enhancer is capable of generating a mild rescue of the *Poxn* null posterior lobe phenotype when fused to Gal4, driving a UAS-Poxn construct (Figure S3L). We observe that the orthologous regulatory region of *D. pseudoobscura* is also capable of generating a similar degree of rescue (Figure S3M). These experiments confirm the ancestral capability of the posterior lobe enhancer region to drive the expression necessary to generate a derived structure, suggesting that an ancestral function of this region was co-opted during the evolution of this novelty. We subsequently considered what this ancestral activity may be.

In the initial screen of the *Poxn* regulatory region (Boll and Noll, 2002), several additional activities of *Poxn* were mapped to a domain overlapping the posterior lobe activity (Figure 2A). As these specificities may represent ancestral functions that were



co-opted as the posterior lobe originated, we examined whether any of these were contained within our reporter fragment. Although many of the described activities were located outside of our reporter construct, strong expression was observed in an embryonic structure—the posterior spiracle (Figure 3A). Indeed, further subdivision of our reporter fragment failed to separate posterior spiracle from posterior lobe activities (Figure S3A). We next evaluated the possibility that the posterior spiracle enhancer of *Poxn* was co-opted during the origination of the posterior lobe.

Shared Topology and Membership of the Posterior Lobe and Spiracle Networks

The posterior spiracle is a larval structure that is connected to the tracheal system, providing gas exchange to the larva (Figure 3D). *Poxn* is expressed in the embryonic region that develops into the posterior spiracle (Figure 3B), and *Poxn* mutants exhibit multiple defects in the spiracle, including transformation of sensory structures (Boll and Noll, 2002) and a shortening of the stigmatophore, an external protuberance that supports the spiracle (Figure 3E). The stigmatophore defect of *Poxn* can be rescued by a transgenic construct containing the posterior lobe and spiracle enhancer fused to a *Poxn* cDNA (Figure 3F).

Figure 2. A Deeply Conserved Enhancer of *Poxn* Is Required for Posterior Lobe Development

(A) Schematic of the *Poxn* locus, displaying a subset of the described enhancer activities (Boll and Noll, 2002) and indicating the relative position of a posterior lobe reporter construct.

(B and C) Accumulation of *Poxn* mRNA during genital development of *D. melanogaster* at (B) 32 hAPF and (C) 48 hAPF.

(D and E) Activity of the *D. melanogaster* posterior lobe reporter at (D) 32 hAPF and (E) 48 hAPF.

(F and G) Expression of *Poxn* in *D. ananassae* showing mRNA accumulation in the region between clasper and lateral plates (F), but not at the site where a lobe would develop (G).

(H and I) Despite the absence of a posterior lobe in *D. ananassae*, the orthologous posterior lobe enhancer region drives expression preceding (H) and during posterior lobe development of *D. melanogaster* (I). CL, clasper; LP, lateral plate; AP, anal plate, PE, penis, PL, posterior lobe. See also Figure S2.

The posterior spiracle is specified during embryogenesis by a network of genes that is activated by the Hox gene *Abdominal-B* (*Abd-B*) (Figure 3G) (Hu and Castelli-Gair, 1999). Intriguingly, genital development also depends upon *Abd-B*, resulting in genital-to-leg transformations in its absence (Estrada and Sánchez-Herrero, 2001).

Considering the apparent parallels between the posterior lobe and the posterior spiracle, we speculated that additional components of the spiracle network

might be active in the developing genitalia. The JAK/STAT pathway plays a critical role in the posterior spiracle network (Lovegrove et al., 2006), and its ligand, encoded by the *unpaired* gene (*upd*, also known as os) (Harrison et al., 1998), is expressed at high levels in the developing posterior lobe (Figure 4A). This pattern is consistent with the activity of a JAK/STAT signaling reporter (Bach et al., 2007), which is expressed at high levels during posterior lobe development (Figure 4B; Figures S4D–S4F). Reduction of JAK/STAT signaling in the genitalia by transgenic RNAi hairpins directed toward the receptor (*dome*), kinase (*hop*), or transcription factor (*Stat92E*) resulted in drastic reductions in the posterior lobe's size compared to a control RNAi hairpin (Figures 4C–4F and 4T). Hence, the major signaling pathway that patterns the posterior spiracle is also active in the novel posterior lobe structure.

We identified three additional top-level transcription factors of the posterior spiracle network that are active during the development of the posterior lobe. Abd-B and Spalt proteins are both deployed in broad domains that include the posterior lobe (Figures 4G and 4H; Figures S4I and S4J), consistent with severe genital defects in *Abd-B* (Estrada and Sánchez-Herrero, 2001; Foronda et al., 2006) and *spalt* mutants (Dong et al., 2003). In contrast, Empty spiracles (Ems), named for its spiracle

Figure 3. The Posterior Lobe Enhancer of *Poxn* Is Active in the Hox-Regulated Network of the Posterior Spiracle

(A) Transgenic embryo bearing the *D. melanogaster* posterior lobe enhancer reporter.

(B) Antibody staining of Poxn protein in the posterior spiracle anlagen of the stage 13 (St13) *D. melanogaster* embryo presented in (A).

(C) Merged image of (A) and (B), showing the *Poxn* enhancer (green) and Poxn protein (magenta).

(D) Scanning electron micrograph of a wild-type (wt) third instar larva, showing the posterior spiracle structure.

(E) The Poxn null mutant posterior spiracle is shorter relative to wild-type.

(F) Rescue of posterior spiracle defects of a *Poxn* mutant by a fragment of the *Poxn* locus containing the lobe and spiracle enhancer fused to a *Poxn* cDNA.
(G) Diagram of the posterior spiracle network, adapted from Hu and Castelli-Gair (1999) and Lovegrove et al. (2006). The addition and placement of *Poxn* and *eya* within this network is based upon data presented in this work.
Arrows in (A)–(F) point to the posterior spiracle. See also Figure S3.

phenotype (Jürgens et al., 1984), is expressed in a restricted genital pattern similar to *Poxn* (Figure 4I; Figure S4K). In summary, five transcription factors required for posterior spiracle development (Abd-B, Poxn, Spalt, Ems, and activated STAT) are deployed in the novel posterior lobe context, suggesting a highly similar trans-regulatory landscape governing these two structures.

While the trans-regulatory landscapes of the lobe and spiracle bear an unexpected resemblance, they also appear to impart a high degree of spatial specificity. Abd-B is restricted to posterior body segments (Celniker et al., 1989), while Poxn, Spalt, and Ems rarely overlap in expression (Dalton et al., 1989; Dambly-Chaudière et al., 1992; Kühnlein et al., 1994). The JAK/ STAT pathway is recurrently deployed during development, but very few tissue settings would include all five factors. We therefore reasoned that downstream genes in the spiracle network might be activated in the developing posterior lobe. To test this possibility, we monitored their expression during genital development. In five genes of this network-engrailed (en), crumbs (crb), Gef64C, Cad86C, and eyes absent (eya)we found corresponding expression within the developing posterior lobe (Figures 4J-4N; Figures S5A-S5E). Hence, a total of at least 10 genes are shared between the 2 networks. We investigated the hierarchal relationship between several of the identified genes by targeting both upper and lower tiers of the network using RNAi hairpins, and measuring downstream effects on gene expression. Reduction of JAK/STAT signaling led to measurable decreases in the expression of Ems, Crb, and Eva (Figures S5F-S5K), while the reduction of crb, Gef64C, or Cad86C did not alter the pattern of Ems expression or JAK/STAT pathway activity measured from the 10X STAT reporter (not shown). Two genes whose expression in the posterior lobe depends on dome have been linked to JAK/STAT activity in the posterior spiracle, crb (Lovegrove et al., 2006) and eya (see below). These results support a shared topology between the two networks.

The sharing of genes between the spiracle and lobe networks may be due to their recent recruitment to posterior lobe development, which would predict that their expression is specific to species that possess this structure. To determine whether the activity of these genes differs between lobed and non-lobed species, we examined their expression in non-lobed species at time points corresponding to stages in which the D. melanogaster lobe emerges. Ems exhibits strong lobe-specific activity that is absent in non-lobed species (Figure S4K); however, both Spalt and Abd-B are widely and strongly expressed in all species tested (Figures S4I and S4J). upd mRNA is weakly present in early genitalia prior to lobe development in both lobed and non-lobed species, but persists and intensifies in D. melanogaster during lobe development (Figure S4A). Downstream spiracle network genes eya, en, crb, Gef64C, and Cad86C are active in several locations within the genitalia, but all exhibit unique lobe-specific expression patterns (Figures S5A-S5E). Thus, of the 10 shared genes that we have discovered, 8 are unique to lobed species during the stages of this structure's emergence.

To confirm that the identified posterior spiracle genes actively participate in posterior lobe development, we targeted *ems*, *crb*, *Gef64C*, *Cad86C*, and *eya* with RNAi hairpins driven by genital drivers. Reduction of *ems*, *Gef64C*, and *Cad86C* significantly reduced the size of the posterior lobe, suggesting that they positively contribute to lobe development, while reduction of *crb* and *eya* significantly increased the size of the lobe compared to a control RNAi hairpin, which may indicate an inhibitory or restrictive role in the lobe's development or expansion (Figures 40–4T). Thus, genes of the spiracle network that are specifically restricted to this novel structure during its development contribute to its construction.

Figure 4. Shared Topology and Membership of the Posterior Lobe and Spiracle Networks

Antibody staining (G–L) and in situ hybridization (A, M, and N) reveal the deployment of several posterior spiracle network genes within the posterior lobe during genital development (arrows).

(A and B) Expression of *upd* mRNA in the developing lobe (A) closely mirrors the activity of a 10XStat92E-GFP reporter (B).

(C–F) Reduction in expression of members of the JAK/STAT signaling pathway *hop* (D), *dome* (E), or *Stat92E* (F) reduces the size of posterior lobe relative to a control (C).

(G-I) The top-tier spiracle network factor Ems (I) is strongly expressed within the developing posterior lobe, while Abd-B (G) and Sal (H) are pre-

Shared Enhancers Underlie the Parallel Topologies of the Lobe and Spiracle Networks

The striking similarity between the posterior lobe and spiracle networks may reflect the convergent evolution of similar network topologies or could result from co-option of the ancestral posterior spiracle network in generating the lobe. To distinguish between co-option and coincidence, we tested additional enhancers of the posterior spiracle network for posterior lobe activity (see Experimental Procedures). In the case of co-option, multiple enhancers of the posterior spiracle network would be active in the posterior lobe, whereas convergence would produce enhancer activities in distinct locations within each shared gene's regulatory region.

The crb gene is deployed in the posterior spiracle through an intronic JAK/STAT responsive enhancer (Lovegrove et al., 2006), which we found to be active in the posterior lobe (Figures 5A, 5G, and 5G'). A recent screen of the regulatory regions of invected (inv) and en identified a posterior spiracle enhancer (Cheng et al., 2014), which consistently drives weak expression during late posterior lobe development (Figures 5B, 5H, and 5H'). We discovered a region of the Gef64C gene that is active in both the posterior spiracle and posterior lobe patterns (Figures 5C, 5I, and 5I'). We also discovered a region of the Cad86C gene that consistently recapitulates a portion of its posterior spiracle expression domain, as well as a lobe-associated pattern that is specific to lobed species (Figures 5D, 5J, and 5J', white arrow). For eya, a new member of the posterior spiracle network identified in this study, we localized an upstream enhancer that recapitulates its genital expression pattern (Figures 5E and 5K'). This enhancer is also active in the outer edge of the larval spiracle's stigmatophore (Figure 5K). While a previously identified posterior spiracle enhancer upstream of ems (Jones and McGinnis, 1993; Figure 5L) lacked activity in the posterior lobe (Figure 5L'), we identified an additional enhancer located just downstream of the transcription unit that is activated in both settings (Figures 5F. 5M. and 5M'). This downstream enhancer of ems recapitulates a previously undescribed activity in the outer edge of the stigmatophore (Figures 5N' and 5P') but is not active in the initial spiracular chamber pattern (Figure 5P). In conclusion, we have identified seven enhancers (Poxn, crumbs, en, Gef64C, Cad86C, eya, and ems) of the posterior lobe network that can be traced to overlapping functions in the posterior spiracle. Given the large size of their respective regulatory regions, we

sent more generally throughout the lateral plate from which the lobe emerges.

(J-N) Downstream spiracle network factors Eya (J) and En (K), as well as terminal differentiation factors Crb (L), *Gef64C* (M), and *Cad86C* (N), are all expressed at specific regions and stages of posterior lobe development.

(O–S) Transgenic RNAi hairpin mediated reduction in expression of spiracle network members *ems* (O), *crb* (P), *Gef64C* (Q), *Cad86C* (R), or *eya* (S) alters the size of posterior lobe compared to a control (shown in C).

(T) Boxplot depicting the relative area of posterior lobes subject to RNAi treatments and normalized to a control. Asterisks denote significant differences (Student's paired t test, *p < 0.05, **p < 0.005). Whiskers denote 1.5 times the interquartile range above and below the first and third quartiles.

Dashed lines mark the position of the developing posterior lobe (A, B, G–N) or demonstrate altered posterior lobe shape (D–F and O–S) compared to a control (C). Arrowhead in (N) identifies a pattern that is not unique to lobed species (Figure S6E). See also Figures S4 and S5.

Figure 5. Co-option of Posterior Spiracle Enhancers to Posterior Lobe Development

(A–F) Schematic diagrams of genomic loci in which an enhancer activated in both the posterior lobe and posterior spiracle were localized (orange boxes). Reporter constructs contain the schematized segment fused to either GFP or Gal4.

(G–M and G'–M') GFP reporter expression driven in transgenic *D. melanogaster* by enhancers for *crb* (G and G'), *en* (H and H'), *Gef64C* (I and I'), *Cad86C* (J and J'), *eya* (K and K'), *ems* upstream (US) enhancer (L and L'), and *ems* DS enhancer (M and M') in the posterior spiracle (G–M) and in the posterior lobe (G'–M'). (N–P) *ems* mRNA is first present at stage 11 in cells that contribute to the spiracular chamber (N), a pattern recapitulated by the *ems* US reporter (O), but not by the *ems* DS reporter (P).

(N'-P') *ems* is also active later during posterior spiracle development around the border of the stigmatophore (N', arrow) and in each embryonic segment, a pattern that is not encoded in the upstream enhancer (O'), but is recapitulated by the *ems* DS reporter (P', arrow). (L') The *ems* US reporter is not expressed within the developing posterior lobe. (J') In addition to a lobe specific pattern (arrow), the *Cad86C* reporter also recapitulates a conserved pattern at the edge of the anal plate (arrowhead).

postulated that the coincidence of lobe and spiracle enhancers would be highly unlikely due to chance alone. Simulations in which we randomized the locations of lobe and spiracle reporter fragments across the full extent of each of the seven loci confirmed an extremely low probability that the observed lobe and spiracle enhancer fragments would overlap by a single nucleotide ($p = 6 \times 10^{-8}$).

The Activation of Enhancers in Both New and Old Contexts Depends on Direct Input from Hox and Signaling Pathway Factors

A hallmark of co-option of regulatory sequences is the use of individual transcription factor binding sites in two or more developmental contexts (Rebeiz et al., 2011). The similarities in lobe and spiracle network topologies and enhancer locations strongly suggested that transcription factor binding sites within posterior spiracle enhancers would be required for posterior lobe function. Therefore, we searched for conserved transcription factor binding sites that could mediate functions common to both networks. Within the Poxn posterior lobe enhancer, we identified instances of high-quality binding sites for STAT and Abd-B, both of which were contained within an 897 bp fragment active in both contexts (Figure S3A). In addition, we identified a high-quality binding site for STAT within a 294-bp interval defined by two overlapping reporters of the eya enhancer that were active in both locations (Figures S6C and S6D). Comparisons to other sequenced Drosophila species revealed that these three sites are highly conserved (Figure 6A), consistent with their potential function in the deeply conserved posterior spiracle structure. Introduction of a 2-bp mutation that is known to disrupt STAT binding (Lovegrove et al., 2006) drastically reduced activity of the Poxn reporter in both the posterior lobe and posterior

Poxn Enhancer eya Enhancer Α STAT BS AbdB BS STAT BS consensus: TTCNNNGAAGCCRTAAA TTCNNNGAA mel GAATTCGTGGAAGCCGTAAAAAG CTCTTCCGAGAATTT sim GAATTCGTGGAAGCCGTAAAAAG CTCTTCCGAGAATTT yak GAATTCGTGGAAGCCGTTAAAAG CTCTTCCGAGAATCG AAATTCGTGGAAGCCGTAAAAAG CTCTTCCGAGAATCG ere GAATTCGTGGAAGCCGTAAAAAG CCCTTCCGAGAATTC tak GATTTCGTGGAAGCCGTAAAAAG GCTTTCCGAGAATTC ele GATTTCGTGGAAGCCGTAAAAAA ACCTTCCGAGAATTC euq GATTTCGTGGAAGCCGTAAAAAG CTTTTCCGAGAATTC ana GATTTCGTGGAAGCCGTAAAACC CCCTTCCAAGAAATG pse wil GATTTCGTGGAAGCCGTAAAATC TGATTCCAAGAAATG GATTTCGTGGAAGCCGTAAAATA CCCTTCCGAGAATTC ari vir GATTTCGGAGAAGCCGTAAAAAA GCCTTCCGAGAAGTG GATTTCGTGGAAGCCGTAAAAAA GCCTTCCGAGAAACG moj STAT mut: AACGTGGAA AACCGAGAA AbdB mut: GCCGCCGA St13 D St13 F St13 St13 St13 E С eya STAT mutant Poxn Poxn eya STAT mutant AbdB mutant enhancer Poxn enhancer 48h C' 48h D' 48h E' 48h Ξž 48h R

Figure 6. Redeployment of *Poxn* and *eya* in the Posterior Lobe Required Ancestral Binding Sites for Abd-B and STAT that Function in the Posterior Spiracle Context

STAT mutant

eya

STAT mutant

AbdB mutant enhancer

(A) Alignment of a Stat92E binding site (purple text) and an Abd-B binding site (green text) of the *Poxn* lobe and spiracle enhancer and a Stat92E binding site (purple text) of the *eya* lobe and spiracle enhancer, showing near perfect conservation among sequenced *Drosophila* species.

(B–D and B'–D') Mutations to two bases in a STAT binding site (C and C') or three bases in an Abd-B binding site (D and D') reduces both posterior spiracle (C and D) and posterior lobe (C' and D') activity compared to the wild-type *Poxn* enhancer (B and B').

(E, E', F, and F') Mutation of two bases in a STAT binding site (F and F') reduces both posterior spiracle (F) and posterior lobe (F') activity compared to the wildtype *eya* enhancer (E and E').

See also Figure S6.

spiracle (Figures 6B, 6B', 6C, and 6C') and similarly eliminated activity of the *eya* enhancer reporter in both settings (Figures 6E, 6E', 6F, and 6F'). Introduction of a 3-bp mutation that disrupts Abd-B binding (Williams et al., 2008) extinguished *Poxn* enhancer activity in the posterior lobe and significantly reduced posterior spiracle expression by 57% (Figures 6D and 6D'). These results demonstrate the co-option of enhancers into a novel setting through the redeployment of pre-existing transcription factor binding sites.

DISCUSSION

Here, we have shown how a GRN underlying a novel structure, the posterior lobe, is composed of components that are active in the embryonic posterior spiracle, an ancestral Hox-regulated structure that was present at the inception of this novelty (Figure 7). These findings confirm previous speculation that network co-option proceeds through the re-use of individual transcription factor binding sites within enhancer sequences (Gao and Davidson, 2008; Monteiro and Podlaha, 2009). Further, our data help calibrate expectations concerning the degree of physical similarity between novel and ancestral structures during co-option events. Below, we briefly discuss how the architecture of the posterior spiracle network may have predisposed it for co-option in the genitalia, and we explore the general implications of our findings with regard to the origins of morphological novelty.

While our results illustrate the downstream consequences of co-option, the upstream causative events await characterization. We suspect that some number of high-level regulators of the posterior spiracle network recently evolved novel genital expression patterns through alterations within their regulatory regions. Currently, Unpaired represents the best candidate upstream factor, as it is positioned near the top of the spiracle network, differs in expression greatly between lobed and nonlobed species (unlike Spalt and Abd-B), and is the only highlevel factor in the spiracle network for which a shared lobe and spiracle enhancer has yet to be identified (Figure 7C). Indeed, a reporter screen of the 30 kb of regulatory DNA immediately surrounding the upd gene identified a posterior spiracle enhancer that is not deployed in the posterior lobe, marking an important point of divergence separating the posterior spiracle and posterior lobe networks (Figures 7D-7F). However, the identification of enhancers controlling upd expression in the posterior lobe will be required to resolve its role in this structure's origination.

The architecture of the posterior spiracle network may have shaped the possible developmental contexts in which it could be co-opted. The Hox factor Abd-B has a deeply conserved role in the insect abdomen and genitalia (Kelsh et al., 1993; Yoder and Carroll, 2006). The top-level factors of the posterior spiracle network depend upon Abd-B for activation in the embryo (Figure 3G) (Hu and Castelli-Gair, 1999). This regulation by Abd-B extends to lower tiers of the network, such as Poxn (Figures 6A, 6D, and 6D'; Table S1). The tight integration of Abd-B with multiple tiers of the posterior spiracle network may have limited this network's re-deployment to posterior body segments that express Abd-B. Indeed, several components of this network (Poxn, ems, upd) are activated early during genital development in the presumptive cleavage furrow separating the lateral plate from the clasper (Figures 2B, S4K, and S4A). This may represent the aftereffect of multiple waves of re-deployment in Abd-B expressing tissues. Examination of additional examples of network co-option at the level of constituent regulatory sequences could reveal general rules that govern and bias network redeployment.

Historically, the identification of co-option events has relied upon comparative analyses of gene co-expression. The first examples of co-option were diagnosed by finding novel gene expression patterns near zones of ancestral function, such as the deployment of the posterior wing patterning circuit within novel butterfly eyespots (Keys, 1999). Subsequently, many examples of co-option have involved educated guesses of the types of networks that contribute to the novelty, such as the

Figure 7. Model Depicting the Co-option of Genes, Enhancers, and Transcription Factor Binding Sites during the Origination of the Novel Posterior Lobe

(A) The posterior spiracle enhancer of Poxn binds Abd-B and phosphorylated STAT in the embryonic posterior spiracle anlagen to activate expression ("ON") (top). In species lacking a posterior lobe. the enhancer is not activated during genital development ("OFF") (middle). The deployment of regulatory factors of the spiracle network during late stages of genital development in lobed species resulted in the activation of the Poxn spiracle enhancer by Abd-B and activated STAT (bottom). (B and C) Summary of Poxn expression (B) and the status of the posterior spiracle network (C) in the three developmental contexts. (C) Expressed genes are shaded in green, while inactive genes are shaded gray. Genes activated by a shared lobe and spiracle enhancer are outlined with red dashes. The yellow dashes surrounding the upd node indicate its activation in the spiracle through an enhancer that lacks lobe activity.

(D) Schematic diagram of the *upd* locus in which a posterior spiracle enhancer was identified (orange box).

(E and F) Reporter construct containing the schematized segment fused to a GFP reporter is active in the posterior spiracle (E), but not in the posterior lobe (F).

(G and H) Illustrated three-dimensional models of the developing posterior spiracle at embryonic stage 13 (G) and the developing posterior lobe (H). Important structural domains for both tissues are identified. The Hox gene Abd-B is expressed in all depicted genital structures and is deployed throughout the entire body segment containing the posterior spiracle. Zones of expression for top-tier factors Spalt (green), Ems (blue), and Cut (red) are shown. The JAK/STAT signaling pathway ligand Unpaired is shown in white, with arrows pointing toward tissues in which the JAK/STAT signaling response has been demonstrated. Downstream network genes Eya (yellow), Poxn, Engrailed, Crumbs, Gef64C, and Cad86C are deployed in distinct portions of both tissues. See also Figure S7.

role of the appendage specification network within beetle horns (Moczek and Nagy, 2005; Moczek et al., 2006) or the sharing of the biomineralization network between adult and larval skeletons of sea urchins (Gao and Davidson, 2008). Our data suggest that tracing the evolutionary origins of individual enhancers provides a less biased path for connecting novelties to their ancestral beginnings, as any of the seven enhancers we have characterized in the posterior lobe would have led us to the spiracle network. Further, this approach is likely to illuminate the underlying cellular mechanisms by which the co-option of a network is translated into a novel developmental outcome.

Rather than generating a serial homolog of the posterior spiracle, the co-option event forming the posterior lobe resulted in an epithelial outgrowth, likely owing to the deployment of only a portion of the spiracle network in the genitalia. This is reflected by the absence of the Cut transcription factor and downstream genes (Figures S7B–S7L) that control the spiracular chamber's development (Hu and Castelli-Gair, 1999). Of the 10 genes we have identified in both networks, 9 are active in the stigmatophore (Figures 7G and 7H), the outer sheath of the posterior spiracle that protrudes from the body through a process that involves convergent extension (Brown and Castelli-Gair Hombría, 2000; Hu and Castelli-Gair, 1999). Collectively, these findings imply that similar morphogenic processes are activated by this shared network in the novel setting of the posterior lobe. We propose that the inspection of enhancers underlying other novel three-dimensional structures may reveal similar networks that have been used over and over again to generate "unique styles of architecture" within developing tissues (Zuckerkandl, 1976).

EXPERIMENTAL PROCEDURES

Fly Strains and Husbandry

All flies were reared on a standard cornmeal medium. Species used in this study were obtained from the University of California, San Diego *Drosophila* Stock Center (*Drosophila biarmipes* #0000-1028.01, *Drosophila ananassae* #0000-1005.01, *Drosophila simulans* #14021-0251.165, *Drosophila pseudoobscura* #0000-1006.01, *Drosophila sechellia* #14021-0248.03, *Drosophila erecta* #14021-0224.01, *Drosophila yakuba* #14021-0261.00). The *Drosophila melanogaster* line used in this study is mutant for *yellow* and *white* (y^1w^1 , Bloomington Stock Center #1495) and was isogenized for eight generations.

Pupal Genital Sample Preparation

To collect developmentally staged genital samples, white prepupae were sorted by sex and incubated at 25° C for 24 to 48 hr. Pupae were cut in half in cold PBS, extricated from the pupal case, and flushed with cold PBS to remove fat bodies and internal organs while preserving the developing genital epithelium. Carcasses were then fixed in PBS with 0.1% Triton-X and 4% paraformadehyde (PBT-fix) at room temperature for 30 min. Samples containing fluorescent reporters were washed three times for 10 min in PBS with 0.1% Triton-X (PBT), then imaged immediately. Samples to be used for in situ hybridization were rinsed twice in methanol and stored in ethanol at -20° C.

Embryo Collection

Embryos were collected from grape agar plates (Genesee Scientific) in egg-lay chambers that were incubated at 25°C for up to 20 hr. Embryos were dechorionated in 50% bleach for 3 min, washed in distilled water, and collected on a nitrile filter. Embryos were then fixed for 20 min in scintillation vials containing PBS, 2% paraformaldehyde, and 50% heptane. The PBS layer was removed from the vial and replaced with an equal amount of methanol. Samples to be used for in situ hybridization were vortexed for 30 s, removed from the methanol layer, rinsed twice in methanol, then stored in ethanol. Samples containing fluorescent reporters or to be used for immunostaining were shaken vigorously by hand for 1 min, rinsed in methanol once, then quickly rinsed in PBT three times to prevent the degradation of GFP and antibody epitopes.

Immunostaining

Embryo and genital samples were incubated overnight at 4°C with primary antibodies diluted in PBT. The following primary antibodies were used: rabbitanti-Poxn 1:100 (Dambly-Chaudière et al., 1992), rabbit anti-Ems 1:200 (Dalton et al., 1989), rabbit anti-Spalt 1:500 (Barrio et al., 1996), mouse anti-Eya 1:100 (Bonini et al., 1997), mouse anti-Crb 1:50 (Tepass and Knust, 1993), mouse anti-Engrailed/Invected 1:500 (Patel et al., 1989), rat anti-E-cadherin 1:100 (antibody DCAD2, Developmental Studies Hybridoma Bank), and mouse anti-Cut 1:100 (antibody 2B10, Developmental Studies Hybridoma Bank), After several washes with PBT to remove unbound primary antibody, samples were incubated overnight in diluted secondary antibody (donkey anti-mouse Alexa 488, and donkey anti-rabbit Alexa 647, both at 1:400 dilution from Molecular Probes, or goat anti-rat Alexa 488 at 1:200 dilution from Molecular Probes) to detect bound primary antibody. Samples were washed in PBT to remove unbound secondary antibody, incubated for 10 min in 50% PBT and 50% glycerol solution, then mounted on glass slides in an 80% glycerol 0.1 M Tris-HCL (pH 8.0) solution.

In Situ Hybridization

The in situ hybridization was performed as previously described in Rebeiz et al. (2009), with the modification that we used an InsituPro VSi robot (Intavis Bioanalytical Instruments). Fixed embryo and genital samples were first dehydrated in a 50% xylenes/50% ethanol solution for 30 min at room temperature. Xylenes were removed by several washes with ethanol before the samples were loaded into the InsituPro VSi. During the automated steps, the samples were washed in methanol, rehydrated in PBT, fixed in PBT-fix, incubated in 1:25,000 proteinase K PBT (from a 10 mg/L stock solution), fixed in PBTfix, and subjected to several washes in hybridization buffer. Samples were probed with digoxygenin riboprobes targeting the coding regions of selected genes (see <u>Supplemental Experimental Procedures</u>, primers for amplifying species-specific mRNA probes) for 18 hr at 65°C. Unbound riboprobe was removed in several subsequent hybridization buffer washes and washed several times in PBT. Samples were removed from the robot and incubated overnight in PBT with 1:6,000 anti-digoxygenin antibody Fab fragments conjugated to alkaline phosphatase (Roche Diagnostics). Alkaline phosphatase staining was then developed for several hours in nitro-blue tetrazolium chloride and 5-bromo-4-chloro-3'-indolyphosphate p-toluidine salt (NBT/BCIP) color development substrate (Promega). Samples were then washed in PBT and mounted on glass slides in an 80% glycerol 0.1 M Tris-HCL (pH 8.0) solution.

Transgenic Constructs

Enhancer elements were cloned using the primers listed in Supplemental Experimental Procedures (primers used for transgenic constructs) and inserted into the vector pS3aG (GFP reporter) or pS3aG4 (Gal4 reporter) using AscI and SbfI restriction sites as previously described (Williams et al., 2008). The primers were designed and sequence conservation was assessed using the GenePalette software tool (Rebeiz and Posakony, 2004). Targeted regions were cloned from genomic DNA purified using the DNeasy Blood & Tissue Kit (QIAGEN). Transcription factor binding site mutations were introduced using overlap extension PCR with mutant primers (see Supplemental Experimental Procedures, primers for generating mutant binding site reporters by overlap extension PCR). All GFP reporters were inserted into the 51D landing site on the second chromosome (Bischof et al., 2007) or the third chromosome 68A4 "attP2" site (Groth et al., 2004) by Rainbow Transgenics. Gal4 insertions depicted in Figure S3 were inserted into the 68E1 landing site on the third chromosome (Bischof et al., 2007). A full list of transgenes and insertions sites is listed in Supplemental Experimental Procedures (transgenic lines analyzed).

The *Poxn* rescue construct depicted in Figure 3F of the main text contains a 7.8-kb genomic fragment containing 3 kb upstream of the Poxn coding unit, including the *Poxn* promoter, and the first three exons and two introns of *Poxn* (which includes the lobe and spiracle enhancer). The remainder of the *Poxn* gene was joined to this construct from a *Poxn* CDNA. This construct ("L2") is identical to the "L1" construct published by Boll and Noll (2002), but differs by the inclusion of 1.5 kb additional sequence upstream of the promoter.

The following GFP and Gal4 reporters were obtained from existing sources. 10XStat92E-GFP reporter was obtained from Erika Bach (Bach et al., 2007). Poxn-Gal4 (construct #13 from Boll and Noll, 2002) and UAS-Poxn was obtained from Werner Boll. *armadillo*-GFP was obtained from the Bloomington *Drosophila* Stock Center (#8556). Several enhancer-GAL4 lines from the Rubin collection (Pfeiffer et al., 2008) were obtained from the Bloomington *Drosophila* Stock Center (BDSC) and are listed in Supplemental Experimental Procedures (transgenic lines analyzed). Transgenic RNAi lines from the Harvard TRiP project include *dome* (#34618), *Stat92E* (#33637) *hop* (#32966), *crb* (#40869), *Cad86C* (#27295), *Gef64C* (#31130), *ems* (#50673), and *eya* (#35725). *mCherry* (#35785), a gene that is not present in the *Drosophila* genome, was used as a control for RNAi experiments. The *salm*-Gal4 driver (#25755) was also obtained from the BDSC.

Microscopy

Adult posterior lobe cuticles and stained in situ hybridization samples were imaged on a Leica M205 stereomicroscope with a $1.6 \times$ objective with the extended multi-focus function. Samples stained with fluorescent antibodies or containing fluorescent reporters were imaged via confocal microscopy at 20× magnification on an Olympus Fluoview 1000 microscope. SEM images of third instar larvae were obtained as previously described by Higashijima et al. (1992).

For each transgenic construct, 3–5 independent lines inserted into the 51D landing site (Bischof et al., 2007) or 68A4 "attP2" landing site (Groth et al., 2004) were derived. A list of reporters and corresponding landing sites are reported in Supplemental Experimental Procedures (transgenic lines analyzed). We compared the relative expression of multiple lines in the genitalia to determine the normal reporter activity of each construct. For quantitative measures, relative fluorescence of the *Poxn* and *eya* posterior lobe enhancers and constructs mutant for STAT and Abd-B sites were determined in both the posterior lobe and posterior spiracle contexts. Mounted genital and embryo samples were imaged at 20× magnification under identical, non-saturating settings uniquely optimized for each sample type. Relative expression within the lobe or spiracle was quantified using ImageJ and assessed using a Student's paired t test.

Simulations of Posterior Lobe and Spiracle Enhancer Co-occurrence

The lengths of shared enhancers and the length of each regulatory region in which these enhancers were embedded were input into an in-house Perl script, CRE-overlap-sim. This program randomizes the location of two equally sized segments of DNA (the size of each reporter fragment tested) across the length of each gene's potential regulatory sequence (the distance from the upstream gene to the gene downstream). For each simulation, the script measures whether the two segments overlapped and counts a successful co-occurrence when all of the input enhancers overlap by the designated number of nucleotides in their respective regulatory regions. A large overlap, which would be expected for co-opted enhancer sequences, will reduce the measured probability of co-occurrence. Our simulations specified a 1 nucleotide overlap, which represents the most permissive and thus most stringent setting possible to detect non-random co-occurrence. 500,000,000 simulations were performed, and the average p value as presented in the main text was calculated.

Identification of Shared and Distinct Posterior Spiracle and Posterior Lobe Enhancers

A combination of comprehensive whole gene surveys and targeted candidate region tests of non-coding regions of genes shared between the two networks was employed to identify co-opted enhancers. In the case of five out of eight of the identified enhancers, multiple constructs inserted in at least two distinct genomic locations were tested for activity. For the whole gene surveys, with the exception of *upd*, we used lines from the Rubin GAL4 collection (Jenett et al., 2012; Pfeiffer et al., 2008), in which non-coding sequences are fused to the GAL4 transcription factor and inserted into the attP2 site on the third chromosome. We supplemented these searches with constructs we generated (see Transgenic Constructs) when necessary. Primers used to amplify reporter constructs are presented in Supplemental Experimental Procedures (primers used for transgenic constructs). We detail the search for each of these enhancers below.

crb

Lovegrove et al. (2006) identified a spiracle enhancer located in the first intron, for which we cloned an identical segment into our reporter system. Additionally, we screened all intronic sequences using the Rubin-Gal4 collection, in which the construct overlapping the Lovegrove fragment uniquely recapitulated lobe expression. We additionally cloned the upstream region of *crb* into our GFP reporter system, and this fragment was not active in the posterior lobe.

en

Cheng et al. (2014) screened the regulatory regions surrounding *engrailed* and *invected*. The "D" enhancer from the intergenic region between *inv* and *en* was shown to specifically recapitulate the posterior spiracle activity of *en*. We reconstituted this enhancer by designing primers to clone the identical segment into our reporter system. This construct drove strong expression similar to endogenous posterior spiracle *en* activity as reported by Cheng et al. (2014) and weak, but consistent, activity within a subset of the posterior lobe, mirroring the levels that appear late during posterior lobe development (Figure S5B).

We screened the upstream region and introns of *eya* using the Rubin Gal4 collection. This screen identified a single posterior lobe activity just upstream of the transcription unit. We further confirmed the activity of this enhancer fragment by inserting it into our GFP reporter system, which showed activity in the posterior lobe as well. Both the GFP reporter and the Rubin Gal4 constructs were expressed in the posterior spiracle. We further refined the size of this regulatory region by testing overlapping fragments of the *D. sechellia eya* enhancer. Two fragments that overlap by 294 bp were active in both spiracle and lobe tissues (Figures S6C, S6C', S6D, and S6D'). The smallest fragment tested was 1,060 bp.

ems

Rubin Gal4 lines existed for nearly all of the ~67-kb region encompassing the non-coding DNA surrounding *ems*. To test a portion of the regulatory region upstream of the *ems* promoter that is not included in the Rubin Gal4 collection, we cloned three additional overlapping regions into our GFP reporter system. We first tested a Rubin Gal4 line that contains the previously identified upstream enhancer for the spiracular chamber (Jones and McGinnis, 1993) (Fig-

ure 5F). This line faithfully reproduced spiracular chamber expression (Figures 5L and 5O), but was not active in the *ems* posterior lobe pattern (Figure 5L'). Screening the other Rubin collection lines of *ems* for genital activity, we identified a fragment just downstream of the transcription unit that drove expression partially recapitulating the lobe expression of *ems*. To determine if this enhancer was indeed distinct from the posterior spiracle activity, we examined its expression in stage 13 embryos and noticed that it was active in the outer stigmatophore (Figures 5M and 5P'), a pattern that recapitulates endogenous *ems* expression (Figure 5N'). We cloned a subfragment of this downstream enhancer into our GFP reporter system, confirming the activity of this segment in the posterior lobe and spiracle. In addition, we cloned the orthologous segment of DNA from *D. ananassae* into our reporter system, demonstrating that a non-lobed species version of *ems* downstream is capable of driving expression within the posterior lobe (Figure S6B').

Gef64C

A survey of the non-coding region of *Gef64C* identified a segment containing several binding sites for genes in the spiracle network, including a high-affinity binding site for Abd-B (Ekker et al., 1994) and two candidate STAT binding sites, all of which were conserved to *D. pseudoobscura*. Fusing this segment of DNA into our reporter system revealed expression in the spiracular chamber of the posterior spiracle, embryonic hindgut, and in several zones in the developing genitalia that recapitulate its endogenous expression (clasper, lobe, anal plate, and hypandrium; Figure 4M). Further truncation of this segment of DNA separated the posterior spiracle and posterior lobe patterns from the other activities, localizing this enhancer to the first intron. This truncation includes the two candidate STAT binding sites, but not the candidate Abd-B binding site (Table S1).

Cad86C

A screen of the non-coding regions surrounding *Cad86C* identified an intronic region near the promoter that included a Spalt site (Barrio et al., 1996), which is conserved to *D. ananassae* (Table S1). We cloned a 3,003-bp segment of DNA that included this region into our reporter system. This reporter consistently recapitulated a portion of the endogenous *Cad86C* activity in the posterior spiracle and embryonic anus (Figure 5J; Figure S5E) and drove expression in the anal plate pattern common to both lobed and non-lobed species (Figure 5J', arrowhead; Figure S5E), as well as the lobe-specific pattern just posterior to the lobe (Figure 5J', arrow; Figure S5E). *upd*

We screened the 30-kb intergenic non-coding DNA between *upd* (also *os*) and its neighboring genes *upd3* and *CG6023* by cloning eight overlapping segments into our reporter system. One reporter directly downstream of *upd* drove expression within the posterior spiracle (Figure 7E), matching the endogenous *upd* pattern (Figure S4G), and none of the tested reporters drove expression within the posterior lobe. The region that drove posterior spiracle expression contains a high-quality match to the Abd-B binding site (Ekker et al., 1994), which is conserved to at least *D. virilis*.

Identification of Predicted Conserved Transcription Factor Binding Sites in Minimal Shared Enhancers

Using the GenePalette Software tool (Rebeiz and Posakony, 2004), we compared the orthologous regions of the shared posterior spiracle and posterior lobe enhancers from *D. melanogaster*, *D. simulans*, *D. yakuba*, *D. biarmipes*, *D. ananassae*, *D. pseudoobscura*, and *D. virilis*. We screened for predicted binding sites for STAT (Yan et al., 1996) and Spalt (Barrio et al., 1996) and for a high-fidelity binding site for Abd-B (Ekker et al., 1994). Putative conserved transcription factor binding sites are listed in Table S1.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, seven figures, and one table and can be found with this article online at http://dx.doi.org/10.1016/j.devcel.2015.08.005.

AUTHOR CONTRIBUTIONS

M.R. and W.J.G. designed the study, performed experiments, and wrote the paper. W.C.J. cloned constructs and established transgenic lines. Y.L. and N.R.D. screened and characterized the *eya* enhancer. S.J.S screened the

upd locus. W.B. and M.N. performed analyses of *Poxn* expression and mutant phenotypes in the posterior spiracle. All authors discussed the results and commented on the manuscript.

ACKNOWLEDGMENTS

The authors thank J.N. Pruitt, T.M. Williams, J. Posakony, and members of the M.R. laboratory for thoughtful discussions and comments on the manuscript. We thank W. McGinnis, R. Barrio, and the Developmental Studies Hybridoma Bank for generously providing antibodies used in this study. *Drosophila* strains were provided by the UCSD *Drosophila* species stock center, the Harvard TRiP, and the Bloomington *Drosophila* stock center. E. Bach, B. Gyeong-Hun, G. Campbell, and A. Kopp provided *Drosophila* stocks. This work was supported by a grant from the NIH (GM107387 to M.R.) and the Alfred P. Sloan Foundation (to M.R.). N.R.D. was supported by an HHMI grant to the University of Pittsburgh.

Received: December 2, 2014 Revised: July 23, 2015 Accepted: August 10, 2015 Published: September 3, 2015

REFERENCES

Abouheif, E. (1999). Establishing homology criteria for regulatory gene networks: prospects and challenges. Novartis Found. Symp. 222, 207–221, discussion 222–225.

Bach, E.A., Ekas, L.A., Ayala-Camargo, A., Flaherty, M.S., Lee, H., Perrimon, N., and Baeg, G.H. (2007). GFP reporters detect the activation of the *Drosophila* JAK/STAT pathway in vivo. Gene Expr. Patterns *7*, 323–331.

Barrio, R., Shea, M.J., Carulli, J., Lipkow, K., Gaul, U., Frommer, G., Schuh, R., Jäckle, H., and Kafatos, F.C. (1996). The spalt-related gene of *Drosophila melanogaster* is a member of an ancient gene family, defined by the adjacent, region-specific homeotic gene spalt. Dev. Genes Evol. 206, 315–325.

Bischof, J., Maeda, R.K., Hediger, M., Karch, F., and Basler, K. (2007). An optimized transgenesis system for Drosophila using germ-line-specific phiC31 integrases. Proc. Natl. Acad. Sci. USA *104*, 3312–3317.

Boll, W., and Noll, M. (2002). The *Drosophila Pox neuro* gene: control of male courtship behavior and fertility as revealed by a complete dissection of all enhancers. Development *129*, 5667–5681.

Bonini, N.M., Bui, Q.T., Gray-Board, G.L., and Warrick, J.M. (1997). The *Drosophila* eyes absent gene directs ectopic eye formation in a pathway conserved between flies and vertebrates. Development *124*, 4819–4826.

Bonner, J.T. (1998). The origins of multicellularity. Integr. Biol. 1, 27-36.

Brown, S., and Castelli-Gair Hombría, J. (2000). *Drosophila* grain encodes a GATA transcription factor required for cell rearrangement during morphogenesis. Development *127*, 4867–4876.

Celniker, S.E., Keelan, D.J., and Lewis, E.B. (1989). The molecular genetics of the bithorax complex of *Drosophila*: characterization of the products of the *Abdominal-B* domain. Genes Dev. *3*, 1424–1436.

Cheng, Y., Brunner, A.L., Kremer, S., DeVido, S.K., Stefaniuk, C.M., and Kassis, J.A. (2014). Co-regulation of invected and engrailed by a complex array of regulatory sequences in *Drosophila*. Dev. Biol. 395, 131–143.

Coyne, J. (1993). The genetics of an isolating mechanism between two sibling species of *Drosophila*. Evolution (N. Y.) 47, 778–788.

Crick, F.H. (1968). The origin of the genetic code. J. Mol. Biol. 38, 367–379.

Dalton, D., Chadwick, R., and McGinnis, W. (1989). Expression and embryonic function of *empty spiracles*: a *Drosophila* homeo box gene with two patterning functions on the anterior-posterior axis of the embryo. Genes Dev. *3* (12A), 1940–1956.

Dambly-Chaudière, C., Jamet, E., Burri, M., Bopp, D., Basler, K., Hafen, E., Dumont, N., Spielmann, P., Ghysen, A., and Noll, M. (1992). The paired box gene *pox neuro*: a determinant of poly-innervated sense organs in *Drosophila*. Cell 69, 159–172.

Davidson, E.H. (2001). Genomic Regulatory Systems: Development and Evolution (Academic Press).

Dong, P.D., Todi, S.V., Eberl, D.F., and Boekhoff-Falk, G. (2003). *Drosophila* spalt/spalt-related mutants exhibit Townes-Brocks' syndrome phenotypes. Proc. Natl. Acad. Sci. USA *100*, 10293–10298.

Eberhard, W.G. (1985). Sexual Selection and Animal Genitalia (Harvard University Press).

Ekker, S.C., Jackson, D.G., von Kessler, D.P., Sun, B.I., Young, K.E., and Beachy, P.A. (1994). The degree of variation in DNA sequence recognition among four *Drosophila* homeotic proteins. EMBO J. *13*, 3551–3560.

Estrada, B., and Sánchez-Herrero, E. (2001). The Hox gene *Abdominal-B* antagonizes appendage development in the genital disc of *Drosophila*. Development *128*, 331–339.

Foronda, D., Estrada, B., de Navas, L., and Sánchez-Herrero, E. (2006). Requirement of *Abdominal-A* and *Abdominal-B* in the developing genitalia of *Drosophila* breaks the posterior downregulation rule. Development *133*, 117–127.

Gao, F., and Davidson, E.H. (2008). Transfer of a large gene regulatory apparatus to a new developmental address in echinoid evolution. Proc. Natl. Acad. Sci. USA *105*, 6091–6096.

Groth, A.C., Fish, M., Nusse, R., and Calos, M.P. (2004). Construction of transgenic *Drosophila* by using the site-specific integrase from phage phiC31. Genetics *166*, 1775–1782.

Harrison, D.A., McCoon, P.E., Binari, R., Gilman, M., and Perrimon, N. (1998). *Drosophila* unpaired encodes a secreted protein that activates the JAK signaling pathway. Genes Dev. *12*, 3252–3263.

Higashijima, S., Michiue, T., Emori, Y., and Saigo, K. (1992). Subtype determination of *Drosophila* embryonic external sensory organs by redundant homeo box genes *BarH1* and *BarH2*. Genes Dev. 6, 1005–1018.

Hu, N., and Castelli-Gair, J. (1999). Study of the posterior spiracles of *Drosophila* as a model to understand the genetic and cellular mechanisms controlling morphogenesis. Dev. Biol. *214*, 197–210.

Jagadeeshan, S., and Singh, R.S. (2006). A time-sequence functional analysis of mating behaviour and genital coupling in *Drosophila*: role of cryptic female choice and male sex-drive in the evolution of male genitalia. J. Evol. Biol. *19*, 1058–1070.

Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D., Cavallaro, A., Hall, D., Jeter, J., et al. (2012). A GAL4-driver line resource for *Drosophila* neurobiology. Cell Rep. 2, 991–1001.

Jones, B., and McGinnis, W. (1993). The regulation of empty spiracles by *Abdominal-B* mediates an abdominal segment identity function. Genes Dev. 7, 229–240.

Jürgens, G., Wieschaus, E., Nüsslein-Volhard, C., and Kluding, H. (1984). Mutations affecting the pattern of the larval cuticle in *Drosophila melanogaster* II. Zygotic loci on the third chromosome. Dev. Biol. *193*, 283–295.

Kelsh, R., Dawson, I., and Akam, M. (1993). An analysis of abdominal-B expression in the locust *Schistocerca gregaria*. Development *117*, 293–305.

Keys, D.N., Lewis, D.L., Selegue, J.E., Pearson, B.J., Goodrich, L.V., Johnson, R.L., Gates, J., Scott, M.P., and Carroll, S.B. (1999). Recruitment of a *hedgehog* regulatory circuit in butterfly eyespot evolution. Science 283, 532–534.

Kopp, A., and True, J.R. (2002). Evolution of male sexual characters in the oriental *Drosophila melanogaster* species group. Evol. Dev. 4, 278–291.

Kühnlein, R.P., Frommer, G., Friedrich, M., Gonzalez-Gaitan, M., Weber, A., Wagner-Bernholz, J.F., Gehring, W.J., Jäckle, H., and Schuh, R. (1994). spalt encodes an evolutionarily conserved zinc finger protein of novel structure which provides homeotic gene function in the head and tail region of the *Drosophila* embryo. EMBO J. *13*, 168–179.

Kuraku, S., Usuda, R., and Kuratani, S. (2005). Comprehensive survey of carapacial ridge-specific genes in turtle implies co-option of some regulatory genes in carapace evolution. Evol. Dev. 7, 3–17.

Lovegrove, B., Simões, S., Rivas, M.L., Sotillos, S., Johnson, K., Knust, E., Jacinto, A., and Hombría, J.C.-G. (2006). Coordinated control of cell

530 Developmental Cell 34, 520–531, September 14, 2015 ©2015 Elsevier Inc.

adhesion, polarity, and cytoskeleton underlies Hox-induced organogenesis in *Drosophila*. Curr. Biol. *16*, 2206–2216.

Moczek, A.P., and Nagy, L.M. (2005). Diverse developmental mechanisms contribute to different levels of diversity in horned beetles. Evol. Dev. 7, 175–185.

Moczek, A.P., and Rose, D.J. (2009). Differential recruitment of limb patterning genes during development and diversification of beetle horns. Proc. Natl. Acad. Sci. USA *106*, 8992–8997.

Moczek, A.P., Rose, D., Sewell, W., and Kesselring, B.R. (2006). Conservation, innovation, and the evolution of horned beetle diversity. Dev. Genes Evol. *216*, 655–665.

Monteiro, A., and Podlaha, O. (2009). Wings, horns, and butterfly eyespots: how do complex traits evolve? PLoS Biol. 7, e37.

Muller, G.B., and Wagner, G.P. (1991). Novelty in evolution: restructuring the concept. Annu. Rev. Ecol. Syst. 22, 229–256.

Orgel, L.E. (1968). Evolution of the genetic apparatus. J. Mol. Biol. 38, 381-393.

Patel, N.H., Martin-Blanco, E., Coleman, K.G., Poole, S.J., Ellis, M.C., Kornberg, T.B., and Goodman, C.S. (1989). Expression of engrailed proteins in arthropods, annelids, and chordates. Cell *58*, 955–968.

Pfeiffer, B.D., Jenett, A., Hammonds, A.S., Ngo, T.T., Misra, S., Murphy, C., Scully, A., Carlson, J.W., Wan, K.H., Laverty, T.R., et al. (2008). Tools for neuroanatomy and neurogenetics in *Drosophila*. Proc. Natl. Acad. Sci. USA *105*, 9715–9720.

Rebeiz, M., and Posakony, J.W. (2004). GenePalette: a universal software tool for genome sequence visualization and analysis. Dev. Biol. 271, 431–438.

Rebeiz, M., Pool, J.E., Kassner, V.A., Aquadro, C.F., and Carroll, S.B. (2009). Stepwise modification of a modular enhancer underlies adaptation in a *Drosophila* population. Science *326*, 1663–1667. Rebeiz, M., Jikomes, N., Kassner, V.A., and Carroll, S.B. (2011). Evolutionary origin of a novel gene expression pattern through co-option of the latent activities of existing regulatory sequences. Proc. Natl. Acad. Sci. USA *108*, 10036–10043.

Robertson, H.M. (1988). Mating asymmetries and phylogeny in the Drosophila melanogaster species complex. Pac. Sci. 42, 72–80.

Sagan, L. (1967). On the origin of mitosing cells. J. Theor. Biol. 14, 255-274.

Small, S., Blair, A., and Levine, M. (1992). Regulation of *even-skipped* stripe 2 in the *Drosophila* embryo. EMBO J. *11*, 4047–4057.

Tepass, U., and Knust, E. (1993). Crumbs and stardust act in a genetic pathway that controls the organization of epithelia in *Drosophila melanogaster*. Dev. Biol. *159*, 311–326.

Williams, T.M., Selegue, J.E., Werner, T., Gompel, N., Kopp, A., and Carroll, S.B. (2008). The regulation and evolution of a genetic switch controlling sexually dimorphic traits in *Drosophila*. Cell *134*, 610–623.

Wittkopp, P.J. (2005). Genomic sources of regulatory variation in *cis* and in *trans*. Cell. Mol. Life Sci. *62*, 1779–1783.

Yan, R., Small, S., Desplan, C., Dearolf, C.R., and Darnell, J.E., Jr. (1996). Identification of a *Stat* gene that functions in Drosophila development. Cell *84*, 421–430.

Yassin, A., and Orgogozo, V. (2013). Coevolution between male and female genitalia in the *Drosophila melanogaster* species subgroup. PLoS ONE *8*, e57158.

Yoder, J.H., and Carroll, S.B. (2006). The evolution of abdominal reduction and the recent origin of distinct *Abdominal-B* transcript classes in Diptera. Evol. Dev. *8*, 241–251.

Zuckerkandl, E. (1976). Programs of gene action and progressive evolution. In Molecular Anthropology, M. Goodman, R. Tashian, and J. Tashian, eds. (Springer), pp. 387–447.

Figure S1, related to Figure 1. A survey of lateral plate morphology and novelty in the oriental lineage. Tree depicting the phylogenetic relationships of selected species in the oriental lineage of *Drosophila*, as well as outgroup species in the *montium*, *ananassae*, and *obscura* groups. The posterior lobe is a cuticular outgrowth unique to the *melanogaster* clade. Species of the *yakuba* clade (*D. yakuba*, *D. santomea*, and *D. teissieri*) exhibit a small outgrowth in a similar position to the posterior lobe that may be homologous to the posterior lobe (Yassin and Orgogozo, 2013). Phylogeny is based upon that of Prud'homme (Prud'homme et al., 2006).

Figure S2, related to Figure 2. Comparison of epithelial morphogenesis and Poxn expression between lobed and non-lobed species. (A-F, A'-F') Confocal images of timed pupal genitalia of an *armadillo*-GFP line highlight changes in epithelial arrangement by monitoring apical cell junctions. (A) At 28 hAPF, the lateral plate (LP) and clasper (CL) form a continuous epithelium. (B) At 32 hAPF, a furrow divides the continuous epithelium into presumptive lateral plate and clasper formations. (C, D) By 40 hAPF, the clasper and lateral plate are completely separated, resembling their adult structures, and the posterior lobe (PL) is visible as a ridge of cells emerging from the lateral plate epithelium. (E, F) by 44 hAPF, the posterior lobe has emerged from the lateral plate, and is in the process of adopting its adult shape. (A'-F') The anal plate (AP) and penis (PE) are adjacent to the lateral plate and clasper epithelial tissue. (G-L) Confocal images of timed D. ananassae pupal genitalia stained with antibody specific for E-cadherin from 28h to 48h demonstrate that these developmental processes are largely conserved except for the development of the posterior lobe. (M) Poxn mRNA is expressed during clasper and lateral plate cleavage in lobed and non-lobed species, but persists in lobed species at the base of the posterior lobe. D. melanogaster and D. simulans develop posterior lobes, D. biarmipes, D. ananassae and D. pseudoobscura do not. All species exhibit Poxn expression during clasper and lateral plate cleavage 28 hAPF to 40h APF, (36 hAPF to 48 hAPF for *D. pseudoobscura*). Only lobed species express *Poxn* during later genital development as the posterior lobe emerges 40 hAPF to 48 hAPF (black arrows, dashed lines). (N) (left) Poxn protein is expressed in a conserved pattern in the posterior spiracles of *D. melanogaster*, *D. biarmipes*, and *D. ananassae*. (right) Poxn protein is highly expressed at the base of the lobe of *D. melanogaster*, but only weak levels of Poxn protein persist, perduring from the earlier phase of expression in the non-lobed species *D. biarmipes* (middle) and *D. ananassae* (bottom).

mel PLE Reporter PLE CD1 Reporter PLE CD2 Reporter PLE CD3 Reporter PLE CD4 Reporter

Figure S3

Figure S3, related to Figure 3. The *Poxn* posterior lobe enhancer (PLE) is inseparable from the posterior spiracle enhancer and pre-dates the evolution of the posterior lobe. (A) Subdivision of the posterior lobe enhancer region. All three activities encoded in this region can be isolated to the smaller cutdown 2 (CD2) truncation construct. Expression designation for each construct is listed (present (+), ectopic, or absent (-)). (B-F) Expression of Poxn reporter constructs in stage 13 embryos, 32 hAPF genital samples (B'-F'), and at 48 hAPF (B"-F"). Compared to the full Poxn PLE reporter, the CD2 region provides the most faithful recapitulation of the larger segment (D-D"). (G-I, G'-I') Reporters of the orthologous regulatory regions of nonlobed species *D. yakuba*, *D. ananassae* and *D. pseudoobscura* exhibit expression during posterior spiracle development at stage 13 (G-I) and posterior lobe development at 48 hAPF (G'-I'). (J-M) *Poxn* mutant flies (J) fail to form a posterior lobe (*Poxn*^{ΔM22-B5}). A UAS-*Poxn* construct (K) is not able to rescue the posterior lobe in the absence of a GAL4 driver (UAS-Poxn, Poxn^{$\Delta M22-B5$}), but when driven by the *Poxn* PLE fused to GAL4 (L), the posterior lobe is partially rescued (*UAS-Poxn*, Poxn^{ΔM22-B5}; mel PLE-GAL4). (M) The Poxn PLE from the non-lobed species D. pseudoobscura is also able to partially rescue the posterior lobe of a *Poxn* mutant (*UAS-Poxn*, *Poxn* $^{\Delta M22-B5}$; pse PLE-GAL4). Dashed lines mark the contour of the $Poxn^{\Delta M22-B5}$ mutant lateral plate for reference.

Figure S4

Figure S4, related to Figure 4. The role of JAK/STAT signaling and top tier posterior spiracle network factors Abdominal-B (Abd-B), Spalt (Sal) and Empty Spiracles (Ems) in posterior lobe development. (A) (left) mRNA of the JAK/STAT ligand upd accumulates in a conserved pattern in the embryonic posterior spiracle at stage 12 (right). In *D. melanogaster* (top), upd mRNA is steadily produced (arrows) both preceding (24h) and during the emergence of the posterior lobe (40-44 h, dashed lines). In the non-lobed species, D. biarmipes (middle), upd mRNA is faintly visible in a region reminiscent of the early phase of *upd* expression in *D. melanogaster*. In D. ananassae (bottom), the second non-lobed species, a similar early pattern of faint upd mRNA accumulation is visible. (B-F) A reporter containing 10 multimerized STAT92E binding sites (Bach et al., 2007) reveals JAK/STAT signaling during cleavage of the clasper from the lateral plate 28 hAPF to 36 hAPF, and during posterior lobe development 40 hAPF to 48 hAPF (arrows), mirroring the expression of upd mRNA. (G) in situ hybridization visualizing mRNA for the JAK/STAT ligand unpaired in a stage 13 embryo. (H) 10XSTAT reporter reveals JAK/STAT signaling is active throughout the stigmatophore portion of the posterior spiracle (PS). (I) Except for the anal plate, Abd-B is expressed throughout the entire genitalia of both lobed (*D. melanogaster*, top) and nonlobed species (D. yakuba, middle; D. erecta, bottom) pupal genitalia at 48 hAPF. (J, K) Sal (J) and Ems (K) expression patterns. Stage 13 embryos (left) and timed pupal genitalia (right) are shown. (J) Spalt is expressed in a conserved pattern within the entire lateral plate of both lobed (D. melanogaster, top) and non-lobed species (D. biarmipes, middle; D. ananassae, bottom). (K) While Ems has a highly conserved pattern in the embryonic posterior spiracle, its genital expression differs greatly between lobed and non-lobed species. In D. melanogaster (top), Ems is highly expressed in the developing posterior lobe. In non-lobed species *D. biarmipes* (middle), and *D.* ananassae (bottom), Ems is only expressed at early stages in the region that will bisect the presumptive clasper and lateral plate, patterns that are visible at 40-44 hAPF.

Figure S5

Figure S5, related to Figure 4. Comparative expression analysis of downstream and terminal spiracle network genes Eya (A), En (B), Crumbs (C), Gef64C (D), and Cad86C (E). (A) (left) Eva exhibits a conserved expression pattern in the outer border of the posterior spiracle of stage 13 embryos. (right) During genital development, Eya is expressed in the lateral plate of lobed and non-lobed species. In the *D. melanogaster* posterior lobe, Eya is repressed in the dorsal portion of the lateral plate and strongly expressed in the ventral portion of the posterior lobe (arrows). (B) (left) en mRNA is expressed in the posterior portion of the posterior spiracle (arrows). (right) En protein is expressed in a highly conserved pattern marking the posterior compartment of the pupal genitalia. In *D. melanogaster*, En is weakly expressed within the posterior lobe (arrows), a pattern that is absent in non-lobed species. (C) Detection of Crumbs protein and *crb* mRNA in *D. melanogaster*, *D. biarmipes*, and *D. ananassae* shows a highly conserved pattern of expression in the posterior spiracle (left), but in the genitalia, expression at high levels in the posterior lobe region is specific to *D. melanogaster* (arrow). (D) *Gef64C* mRNA accumulates in a highly conserved pattern in the developing spiracle (left), but much like Crumbs, is specifically expressed in the posterior lobe of *D. melanogaster*. (E) mRNA pattern for *Cad86C* is conserved in the posterior spiracle (left) but lobe-associated expression is unique to the lobebearing species *D. melanogaster* (top, arrow). Arrowheads mark an anal plate associated pattern that appears in both lobed and non-lobed species. (F-K). RNAi hairpins targeted at JAK/STAT signaling component *dome* reduces the expression of Crb (G), Ems (I) and Eya (K) compared to controls (F, H, J). Values in panels F-K denote the staining intensity expressed as a percentage normalized to a control ± S.E.M.

Figure S6, related to Figure 5. A conserved enhancer of *empty spiracles (ems)* was co-opted to the posterior lobe, and the spiracle and lobe activities of an *eyes absent (eya)* enhancer are inseparable. (A-B, A'-B') Orthologous regions of the *ems DS* enhancer from lobed species *D. melanogaster* (A, A') and non-lobed species D. ananassae (B, B') both drive expression in the posterior spiracle (A, B) and posterior lobe (A', B'). (C-D, C'-D') Two overlapping subfragments of the *eya* spiracle/lobe enhancer cloned from *D. sechellia*, a lobed species, maintain both posterior lobe (C'-D') and posterior spiracle (C-D) activity (arrows).

Figure S7, related to Figure 7. Spiracular chamber genes are not expressed during posterior lobe development. (A-D) Cut antibody staining reveals expression in the spiracle (A) as previously reported (Hu and Castelli-Gair, 1999), but in the developing genitalia, expression is limited to developing sensory organs (B-D). (E-H) The spiracular chamber gene *Cad88C* is expressed at high levels in the spiracle (e), but is absent during genital development (F-H). (I-L) A similar absence of *Cad96Cb* during posterior lobe development is observed. Dashed lines mark the position of the posterior lobe.

Minimum enhancer	Predicted Binding Site	Extent of conservation	Position	Sequence
Poxn CD2 enhancer	high fidelity AbdB	to D. vir	2R:15831429	GCCGTAAA
	STAT	to D. vir	2R:15831437	TTCGTGGAA
crb enhancer	weak STAT	to D. vir	3R:24297639	TTCGTTTGAA
	weak STAT	to D. pse (strong STAT in D. vir)	3R:24297680	TTCAGGGGAA
		,		
Gef64C CD4 enhancer	weak STAT	to D. pse	3L:4696998	TTCCGTGGAA
	weak STAT	to D. vir	3L:4697343	TTCTGTTGAA
Cad86C enhancer	Spalt	to D. pse	3R:10828636	TTATGTAAT
eya CD6 enhancer	STAT	to D. vir	2L:6550780	TTCCGAGAA
ems DS enhancer	high fidelity AbdB	to D. ana	3R:13906140	TTTATGGC
	Spalt	to D. pse	3R:13905120	TTATGAAAT
	STAT	to D. vir	3R:13905989	TTCTCGGAA
	STAT	to D. vir	3R:13906567	TTCCTGGAA

Table S1, related to Figure 7. Putative conserved STAT, Spalt and high fidelity Abd-B binding sites in minimal enhancers shared between posterior spiracle and posterior lobe networks. Predicted binding sites in minimal co-opted enhancers of *D.melanogaster* were compared to the orthologous regions from *D. simulans, D. yakuba, D. biarmipes, D. ananassae (D. ana), D. pseudoobscura (D. pse)* and *D. virilis (D. vir)*, only sites conserved to *D. ana, D. pse*, and *D. vir* are listed. Consensus binding sites are as follows, STAT: TTCNNNGAA, weak STAT: TTCNNNNGAA, Spalt: TTATGWAMT, high fidelity Abd-B: TTTAYGGC.

Supplemental Experimental Procedures

Gene	Species	Forward Primer	Reverse Primer
poxn	D. melanogaster	ACCGTGGTGAAGAAGGATCATCC	taatacgactcactataggCAGATCAAAACTGGGTCAGTGG
poxn	D. biarmipes	atttaggtgacactatagaGAGGAGAACAGCGGCATGTTG	taatacgactcactataggGATTCCACAGCCAGTGCTTGTG
poxn	D. ananassae	atttaggtgacactatagaTTCCTTACTAGATTTTCACTGTTC	taatacgactcactataggTCGATGGAGCTCTCCGAC
poxn	D. pseudoobscura	atttaggtgacactatagaTTCCTTACTAGATTTTCACTGTTC	taatacgactcactataggCGCTTTGATGGATGTCGTCGTGG
ems	D. melanogaster	CGACGAACTGATCGACATGGAGATG	taatacgactcactataggTATAGTTGGTGGTGTCTAGCCTAGG
upd	D. melanogaster	TTCTAGTCACATAAGAGCAACCGC	taatacgactcactatagggagaTCAAGCACTATATCACAGAT
upd	D. biarmipes	TTCTAGTCACATAAGAGCAACCGC	taatacgactcactatagggagaTCAAGCACTATATCACAGAT
upd	D. ananassae	GTAGCTTAAGTAAATTATTTGATTG	taatacgactcactatagggagaGCGGTTGCTCTTATGTGACTAGAA
Cad86C	D. melanogaster	ACAACAACGGCACGTTCGAGATCAG	taatacgactcactataggCATCACTTCGCGATCGAAGCCATGC
Cad86C	D. biarmipes	ACAGCCAAAGACGAYCTTCATC	taatacgactcactataggTDATCTGCTTGCCATCYGGYTGCTC
Cad86C	D. ananassae	ACAGCCAAAGACGAYCTTCATC	taatacgactcactataggTDATCTGCTTGCCATCYGGYTGCTC
Cad88C	D. melanogaster	TGCCATAGTGCTAACGCTGACTGAC	taatacgactcactataggATCCTCCAGATCCTTTACCTTCACC
Cad96Cb	D. melanogaster	CCATTCAGTACACGATAGTCCAGTC	taatacgactcactataggCATCTTCTCGTAGTCGAGTGGCTTG
crb	D. melanogaster	GACAACGGCTATAACCACCTGATCG	taatacgactcactataggATCATCGGACACCTCACCAGGTAAC
crb	D. ananassae	CAGACGAACCCCTGCCTGAACAATG	taatacgactcactataggACCAAATATGCTTGCCGCRCGATCC
Gef64C	D. melanogaster	GAGACGGAGCTCTTGAAGATTCTTC	taatacgactcactataggGAAATCGAAGAGCTCGTAGTTGTGG
Gef64C	D. biarmipes	CGCGATTATGACGATGATGACGAGC	taatacgactcactataggTTYATGCGCAGCGCCATTGTGTCC
Gef64C	D. ananassae	CGCGATTATGACGATGATGACGAGC	taatacgactcactataggTTYATGCGCAGCGCCATTGTGTCC
eya	D. melanogaster	AAGACCACGCCCACGGGYAAGWC	taatacgactcactataggTGACATCGTCGATGTGCACCTGGTC
eya	D. biarmipes	AAGACCACGCCCACGGGYAAGWC	taatacgactcactataggTGACATCGTCGATGTGCACCTGGTC
eya	D. ananassae	AAGACCACGCCCACGGGYAAGWC	taatacgactcactataggTGACATCGTCGATGTGCACCTGGTC
en	D. melanogaster	TSTGCAAGGCGGTCTCSCAGATYGG	taatacgactcactataggTGGTKGTGGATCCCGTCTCSGARCG
en	D. biarmipes	TSTGCAAGGCGGTCTCSCAGATYGG	taatacgactcactataggTGGTKGTGGATCCCGTCTCSGARCG
en	D. ananassae	TSTGCAAGGCGGTCTCSCAGATYGG	taatacgactcactataggTGGTKGTGGATCCCGTCTCSGARCG
Gal4	S cerevisiae	atttaggtgacactatagaTCACAGTGTGCAATCCCATTACCGC	taatacqactcactataqqGGACCGTTGCTACTGTTAGTGAAAG

Primers for amplifying species-specific mRNA probes. Lowercase letters represent sequences for the SP6 (Forward primer) and T7 (Reverse primer) RNA polymerases used for probe synthesis.

Construct	Forward Primer	Reverse Primer
mel poxn PLE	TTCCGggcgcgccTCGGTGGCTTAACACGCGCATT	TTGCCcctgcaggATCGCTGATTCCATGGCCCAGT
yak poxn PLE	TTCCGggcgcgccTTAACGGCCAGCACGAGTTTCC	TTGCCcctgcaggCGTTTTGTTCGAGCGAGTGCAG
ana poxn PLE	TTCCGggcgcgccTTAACGGCCAGCACGAGTTTCC	TTGCCcctgcaggCGTTTTGTTCGAGCGAGTGCAG
pse poxn PLE	TTCCGggcgcgccTTAACGGCCAGCACGAGTTTCC	TTGCCcctgcaggCGTTTTGTTCGAGCGAGTGCAG
mel poxn PLE G4	TTCCGggcgcgccTCGGTGGCTTAACACGCGCATT	TTGCCcctgcaggATCGCTGATTCCATGGCCCAGT
pse poxn PLE G4	TTCCGggcgcgccTTAACGGCCAGCACGAGTTTCC	TTGCCcctgcaggCGTTTTGTTCGAGCGAGTGCAG
mel poxn PLE CD1	TTCCGggcgcgccTCGGTGGCTTAACACGCGCATT	TTGCCcctgcaggGTGATTCGATACGATCCGATGC
mel poxn PLE CD2	TTCCGggcgcgccACTGGTCACTGGACATGGCCAT	TTGCCcctgcaggCCTAAGCCTCCCAATAGAGCGA
mel poxn PLE CD3	TTCCGggcgcgccGTTGATCACATTTCAGCCATGC	TTGCCcctgcaggCCATGGGAAACCAGAAGCTGG
mel poxn PLE CD4	TTCCGggcgcgccTCGCTCTATTGGGAGGCTTAGG	TTGCCcctgcaggATCGCTGATTCCATGGCCCAGT
enhancer	TTCCGggcgcgccTAAACGCAGTACGTGGGCGTTGCAC	TTGCCcctgcaggTGTTGTCCGCGGCTCAATTGTTTGG
Gef64C Intron enhancer	TTGCCcctgcaggTTGCCcctgcaggCAACCAACCCACTTGTGAAGGACTG	TTGCCcctgcaggCACGATTCTTCTTCGCCGAAGAACG
Gef64C Intron CD1	TTGCCcctgcaggCACGATTCTTCTTCGCCGAAGAACG	TTCCGggcgcgccCTGCAAGAGGGAGCTCGTCTACAAG
Gef64C Intron CD2	TTGCCcctgcaggCCCAATGAGCATAAAGCTAATGAGG	TTCCGggcgcgccGCAATCCTCAGACACTTAGTCACCG
Gef64C Intron CD3	TTGCCcctgcaggGAAGATCTCCGGTCCGAAGATGTCC	TTCCGggcgcgccCGACAGCTTCCAATTCAACGCGCTC
Gef64C Intron CD4	TTGCCcctgcaggTTGCCcctgcaggCAACCAACCCACTTGTGAAGGACTG	TTCCGggcgcgccAACAGGTCAAGTGCCGCTTGTCTAC
engrailed D enhancer	TTCCGggcgcgccGAATTCGACGCTTAACTAATGATGC	TTGCCcctgcaggGAATTCGCTTGGCTCACACTGAAAC
Cad86C intron enhnacer	TTCCGggcgcgccGCGAAGACAGATACCGAGATGGTC	TTGCCcctgcaggATTAAAGACGTGCTGGACGCGGAAG
eyes absent PLE	ccgggcgaattcgccggcgcgccTCCTAATTCCATCCGACTTTAAGC	cggttgcgatcgcttcctgcaggTGACTTGTTAAATGGGTGTTCC
eyes absent PLE CD5	TTGCCcctgcaggGGAAGGTGGTGGTGGGTTTTTAAGG	TTCCGggcgcgccTACATGACAAAGCTGCTGGGGATGC
eyes absent PLE CD6	TTGCCcctgcaggTCGACCCATCATCATCTTGATGAGC	TTCCGggcgcgccAGGGGTTGGGTAGCTTAAGTTGTCG
ana ems DS enhancer	TTGCCcctgcaggGWRTTYGTCCCACTGTGTGACAGWG	TTCCGggcgcgccGAGATGCATATCAACAATTAGGACGC
ems US hole 1	TTCCGggcgcgccGACGCAAGTCATTCGGGATATGGG	TTGCCcctgcaggCGTGTGGAGCTTGTAAATGACTCAG
ems US hole 2	TTCCGggcgcgccATCATTTACGAAGAAAGCGAGCCGG	TTGCCcctgcaggACGGTAGCCGTCTATCAGATCAGTG
ems US hole 3	TTCCGggcgcgccTCTGAAGAGTTCTCGTCAAGCAGGC	TTGCCcctgcaggCTCACACTGTATCGCCTCCGCTTAG
upd US 1	TTGCCcctgcaggTGCTATCACTGTTCCTCCCTGACTAG	TTCCGggcgcgccCTGAGAAATGGGAAACTCACACCTC
upd US 2	TTGCCcctgcaggCTTTCGAGGGCTTGCACAATTGACG	TTCCGggcgcgccCATACGCGTACCACCATACTCACTG
upd DS1	TTCCGggcgcgccTCCTGGCGCCATATCAATTACACTC	TTGCCcctgcaggTCGGATGCAAAGTATGTGCACATGG
upd DS2	TTCCGggcgcgccCTCTTGACCTTTTGCGGCTATTTGG	TTGCCcctgcaggTCCAGTACACATATCTTCGCGTAGG
upd DS3	TTCCGggcgcgccCTTTCGTCGTCAGCTCGTCAGTTTG	TTGCCcctgcaggTCATCTCATCTCAGCTCCAGACACC
upd DS4	TTCCGggcgcgccGTTCACCTTGTTTATGGACTCGCTG	TTGCCcctgcaggAGACAGAGAGAGGGGGATCAGAAACC
upd DS5	TTCCGggcgcgccATGCATCAATTAGCTCCCACTGAGC	TTGCCcctgcaggGTAGCGGTAGCAAAAGGCTACTAAC
upd DS6	TTCCGggcgcgccGAGATGCTGTGCCGGTGATTATGAC	TTGCCcctgcaggACCGACATATGACTAAGCCAGCAGC

Primers used for transgenic constructs. Lowercase letters represent restriction sites for *AscI* (Forward primers) and *Sbf I* (Reverse primers) used for cloning

Mutation	Forward Primer	Reverse Primer			
mel poxn PLE STAT mutant	CGCCGTCCGAAaaCGTGGAAGCCG	CGGCTTCCACGttTTCGGACGGCGTCCAGGCCC			
mel poxn PLE Abd-B mutant	CGAATTCGTGGAAGCCGccgAAAGTCTTCGGGGAGTG	CACTCCCCGAAGACTTTcggCGGCTTCCACGAATTCG			
mel eya STAT mutant	CTGCAGCTCaaCCGAGAATTTGGTACGAG	CTCGTACCAAATTCTCGGttGAGCTGCAG			
D I I					

Primers for generating mutant binding site reporters by overlap extension PCR. Lowercase letters represent altered bases introduced into the mutant construct to disrupt the binding sites.

Papartar Nama	Enocioc	5100 CED	68A4 (attP2),	68A4 (attP2),	6951 6514
	Species	X (Fig. 2D, 2E, 6B, 6B', S3B,	ULL.	Gait	
Poxn posterior lobe enhancer	mei	S3B", S3B")			X (Fig S31')
Poxn posterior lobe enhancer	уак	X (Fig. 53G, 53G)			X
Poxn posterior lobe enhancer	ana	X (Fig. 2H, 2I, S3H, S3H')			X
Poxn posterior lobe enhancer	pse	X (Fig. S31, S31')			X (Fig S3M')
Poxn posterior lobe enhancer CD1	mel	X (Fig. S3C, S3C', S3C'')			
Poxn posterior lobe enhancer CD2	mel	X (Fig. S3D, S3D', S3D'')			
Poxn posterior lobe enhancer CD3	mel	X (Fig. S3E, S3E', S3E'')			
Poxn posterior lobe enhancer CD4	mel	X (Fig. S3F, S3F', S3F'')			
Poxn enhancer STAT mutant	mel	X (Fig. 6C, 6C', S3J)			
Poxn enhancer AbdB mutant	mel	X (Fig. 6D, 6D', S3K)			
			X (Fig. 5G,		
crb enhancer	mel	X	5G')	48944	
crb US	mel	X			
crb Rubin Gal4 line with no activity	mel			48851	
crb Rubin Gal4 line with no activity	mel			48877	
crb Rubin Gal4 line with no activity	mel			48918	
crb Rubin Gal4 line with no activity	mel			45455	
en D enhancer	mel	X (Fig. 5H, 5H')			
Gef64C Introp \pm Evon 1	mel	У	Y		
	mol	×	^		
	mer	Χ			
Ger64C IE CD2	mel	X			
Gef64C IE CD3	mel	X			
Gef64C IE CD4	mel	X (Fig. 5I, 5I')			
			V (F)		
Cad86C enhancer	mel		X (Fig.		
	mei		(در,در		
			X (Fig.	48893 (Fig.	
eya enhancer	mel	Х	6E,6E')	5K,5K')	
-,			X (Fig.	- ,- ,	
eya enhancer STAT mutant	mel		6F,6F')		
eya enhancer CD5	sech	X (Fig. S6C, S6C')			
eva enhancer CD6	sech	X (Fig. S6D, S6D')			
	mel	X (1.9. 000 / 000 /		48881	
ava Bubin Cald line with no activity	mel			40001	
B Li G IA II III III ACUVILY	mer			47690	
eya Rubin Gal4 line with no activity	mei			48891	
eya Rubin Gal4 line with no activity	mel			45837	
eya Rubin Gal4 line with no activity	mel			48897	
eya Rubin Gal4 line with no activity	mel			49292	
eya Rubin Gal4 line with no activity	mel			48930	
				47890 (Fig. 5L,	
ems US enhancer	mel			5L', 50, 50')	
				40525 (FIG. 5M), 5M' 5D 5D' 66A	
ems DS enhancer	mel			S6A')	
			X (Fig. S6B,	,	
ems DS enhancer	ana		S6B')		
ems DS enhancer CD1+2	mel	X			
ems US hole 1	mel	Х			
ems US hole 2	mel	Х			
ems US hole 3	mel	Х			
ems Rubin Gal4 line with no activity	mel			48387	
ems Rubin Gal4 line with no activity	mel			46846	
ems Rubin Gal4 line with no activity	mel			40510	
ems Rubin Gald line with no activity	mel			40310	
ems Rubin Gal4 line with no activity	mer			41318	
erns Rubin Gai4 line with no activity	mei			46847	
ems Rubin Gal4 line with no activity	mel			46848	
ems Rubin Gal4 line with no activity	mel			47828	
ems Rubin Gal4 line with no activity	mel			48397	
ems Rubin Gal4 line with no activity	mel			40522	
ems Rubin Gal4 line with no activity	mel			49423	
ems Rubin Gal4 line with no activity	mel			47981	
ems Rubin Gal4 line with no activity	mel			49424	
ems Rubin Gal4 line with no activity	mel			46856	
ems Rubin Gald line with no activity	mel			46857	
ame Rubin Cald line with an article	mel			40007	
erns Rubin Gal4 line with no activity	mei			46858	
ems Rubin Gal4 line with no activity	mel			40528	
ems Rubin Gal4 line with no activity	mel			46865	
ems Rubin Gal4 line with no activity	mel			40529	
upd enhancer DS1	mel	Х			

upd enhancer DS2	mel	Х		
upd enhancer DS3	mel	х		
upd enhancer DS4	mel	х		
upd enhancer DS5	mel	х		
upd enhancerDS6	mel	X (Fig. 7E, 7F)		
upd enhancer US1	mel	X		
upd enhancer US2	mel	Х		

Transgenic lines analyzed. For each construct generated, the species from which it was cloned is listed, as well as the insertion site, and type of reporter (GFP or GAL4). For GAL4 constructs from the Rubin collection, the Bloomington Drosophila Stock Center stock number is listed. For each transgenic line that was presented in a figure, the panel number is provided.

Supplemental References

Bach, E.A., Ekas, L.A., Ayala-Camargo, A., Flaherty, M.S., Lee, H., Perrimon, N., and Baeg, G.H. (2007). GFP reporters detect the activation of the Drosophila JAK/STAT pathway in vivo. Gene Expr Patterns 7, 323–331.

Hu, N., and Castelli-Gair, J. (1999). Study of the posterior spiracles of Drosophila as a model to understand the genetic and cellular mechanisms controlling morphogenesis. Dev. Biol. 214, 197–210.

Prud'homme, B., Gompel, N., Rokas, A., Kassner, V.A., Williams, T.M., Yeh, S.D., True, J.R., and Carroll, S.B. (2006). Repeated morphological evolution through cis-regulatory changes in a pleiotropic gene. Nature 440, 1050–1053.

Yassin, A., and Orgogozo, V. (2013). Coevolution between Male and Female Genitalia in the Drosophila melanogaster Species Subgroup. PLoS One 8, e57158.